

# Care CEUs

## Mental Health Disorders: Identification and Treatment

**1. Major depressive disorder may refer to a disorder characterized by depression that occurs most days of the week for a period of 2 weeks or longer, leading to clinically significant distress or impairment in social, occupational, or other important areas of functioning.**

- A. True**
  - B. False**
- 

**2. A 28-year-old male patient has questions regarding major depressive disorder. Which of the following educational points should be expressed to the patient?**

- A. Anhedonia is a symptom of major depressive disorder.**
  - B. Fatigue is not typically a symptom of major depressive disorder.**
  - C. Major depressive disorder does not typically cause clinically significant distress or impairment in social functioning.**
  - D. In order to be diagnosed with major depressive disorder, the symptoms of major depressive disorder must result from another medical condition.**
- 

**3. A fellow health care professional has questions regarding generalized anxiety disorder. Which of the following informational points of interest should be communicated to the health care professional regarding generalized anxiety disorder?**

- A. Irritability is not a symptom of generalized anxiety disorder.**
  - B. Mania is a symptom of a generalized anxiety disorder.**
  - C. In order for an individual to be diagnosed with generalized anxiety disorder, he or she must exhibit symptoms for at least 2 weeks.**
  - D. The anxiety, worry, or physical symptoms associated with generalized anxiety disorder typically cause clinically significant distress or impairment in social, occupational, or other important areas of functioning.**
- 

**4. Which of the following statements is most accurate?**

- A. Bipolar I disorder is characterized by the occurrence of 1 or more manic or mixed episodes.**
- B. Bipolar I disorder is characterized by the occurrence of 1 or more hypomanic or mixed episodes.**

- C. Bipolar 2 disorder is characterized by the occurrence of 1 or more manic or mixed episodes.
- D. Bipolar 2 disorder is characterized by the occurrence of 1 or more manic episodes.
- 

5. Which of the following statements regarding the DSM-5 criteria for bipolar II disorder is most accurate?

- A. In order to be diagnosed with bipolar II disorder, an individual must have experienced at least 1 major depressive episode.
- B. In order to be diagnosed with bipolar II disorder, an individual must have experienced at least 1 major depressive episode before the age of 65.
- C. In order to be diagnosed with bipolar II disorder, an individual must have experienced at least 2 major depressive episodes.
- D. In order to be diagnosed with bipolar II disorder, an individual must have experienced at least 2 major depressive episodes before the age of 65.
- 

6. Interpersonal therapy is a form of long-term psychotherapy, which typically lasts several years.

- A. True
- B. False
- 

7. Which of the following statements regarding electroconvulsive therapy (ECT) is most accurate?

- A. ECT is typically used as first line therapy to treat mental health disorders for patients < 25 years of age.
- B. ECT is typically used as first line therapy to treat mental health disorders for patients > 25 years of age.
- C. ECT is typically used as first line therapy to treat mental health disorders for patients > 65 years of age.
- D. ECT is often used for patients where other therapeutic options have failed.
- 

8. Which of the following statements is most accurate?

- A. 20 mg/day is the maximum recommended dose of Celexa for patients who are greater than 60 years of age.
- B. 30 mg/day is the maximum recommended dose of Celexa for patients who are greater than 60 years of age.
- C. 40 mg/day is the maximum recommended dose of Celexa for patients who are greater than 60 years of age.
- D. 60 mg/day is the maximum recommended dose of Celexa for patients who are greater than 60 years of age.
-

**9. A fellow health care professional has questions regarding Prozac. Which of the following points of interest should be expressed to the health care professional.**

- A. A typical adult starting dose for Prozac is 20 mg daily.**
  - B. A typical adult starting dose for Prozac is 80 mg daily.**
  - C. Prozac may be taken with monoamine oxidase inhibitors (MAOIs).**
  - D. Prozac may be taken with monoamine oxidase inhibitors (MAOIs) if the patient is < 65 years of age.**
- 

**10. Lexapro is not approved for use in pediatric patients less than 12 years of age.**

- A. True**
  - B. False**
- 

**11. Which of the following statements is most accurate?**

- A. Ativan is not recommended for use in patients with primary depressive disorder or psychosis.**
  - B. Ativan is indicated for the treatment of primary depressive disorder.**
  - C. Ativan is indicated for the treatment of psychosis.**
  - D. Ativan may be used to treat primary depressive disorder in patients over the age of 65.**
- 

**12. Warnings associated with Valium include the following: concomitant use of benzodiazepines, including Valium, and opioids may result in profound sedation, respiratory depression, coma, and death.**

- A. True**
  - B. False**
- 

**13. Which of the following statements is most accurate?**

- A. Lithium toxicity is not closely related to serum lithium concentrations.**
  - B. Lithium toxicity is only related to serum lithium concentrations when doses of lithium exceed 300 mg daily.**
  - C. Lithium toxicity is closely related to serum lithium concentrations, however lithium toxicity cannot occur at doses close to therapeutic concentrations.**
  - D. Lithium toxicity is closely related to serum lithium concentrations, and can occur at doses close to therapeutic concentrations.**
- 

**14. Which of the following statements is most accurate?**

- A. Lamictal is indicated for the maintenance treatment of bipolar I disorder in patients aged 16 years or younger.**
  - B. Lamictal is indicated for the maintenance treatment of bipolar I disorder in patients aged 18 years or younger.**
  - C. Lamictal is not typically associated with Stevens-Johnson syndrome.**
  - D. Lamictal carries warnings regarding Stevens-Johnson syndrome.**
- 

**15. Breastfeeding is not recommended while taking Zyprexa.**

- A. True**
  - B. False**
- 

**Copyright © 2024 Care CEUs**

**Visit us at <https://www.careceus.com>**