

Care CEUs

COVID-19: An Update for 2021

1. A patient has questions regarding COVID-19. Which of the following educational points should be expressed to the patient?

- A. COVID-19 can spread through person-to-person contact.
 - B. COVID-19 is primarily transmitted through the air.
 - C. COVID-19 is only transmitted by contact with infected blood.
 - D. COVID-19 is only transmitted via contaminated food.
-

2. According to the course, how long can the COVID-19 virus live on surfaces?

- A. Up to 7 days
 - B. Up to 10 days
 - C. Up to 14 days
 - D. Up to 28 days
-

3. According to the course, what is the suggested incubation period for COVID-19?

- A. 1 - 14 days
 - B. 14 - 18 days
 - C. 18 - 20 days
 - D. 20 - 30 days
-

4. According to the course, which of the following may be a symptom of COVID-19?

- A. Seizures
 - B. Weight gain
 - C. Fever
 - D. Rash
-

5. A health care professional has questions regarding antigen tests for COVID-19. Which of the following informational points of interest should be communicated to the health care professional?

- A. Antigen tests should not be used on individuals over the age of 65.
 - B. Antigen tests should not be used on individuals under the age of 65.
 - C. Antigen tests may not be as sensitive as NAATs, and the results of an antigen test may need to be confirmed with a NAAT.
 - D. Antigen tests are more sensitive than NAATs.
-

6. According to the course, what is the recommended dose for the Pfizer-BioNTech COVID-19 vaccine?

- A. 0.2 mL
 - B. 0.3 mL
 - C. 0.4 mL
 - D. 0.5 mL
-

7. A 48-year-old female patient has questions regarding the Moderna COVID-19 vaccine. Which of the following educational points should be expressed to the patient?

- A. The vaccine should be administered in 2-dose series separated by 18 days.
 - B. The vaccine should be administered in 2-dose series separated by 28 days.
 - C. The vaccine cannot be offered to individuals that have a history of prior symptomatic or asymptomatic COVID-19 virus infection.
 - D. The vaccine cannot be offered to individuals, over the age of 40, that have a history of prior symptomatic or asymptomatic COVID-19 virus infection.
-

8. A health care professional has questions regarding bamlanivimab. Which of the following informational points of interest should be communicated to the health care professional?

- A. Bamlanivimab should be considered the standard of care for the treatment of patients with COVID-19.
 - B. Bamlanivimab should be considered the standard of care for the treatment of patients over the age of 18 with COVID-19.
 - C. Bamlanivimab must be administered by IV infusion.
 - D. Bamlanivimab must be administered by IM injection.
-

9. What is the recommended adult Day 1 dose of remdesivir for the treatment of COVID-19?

- A. 100 mg
 - B. 200 mg
 - C. 400 mg
 - D. 500 mg
-

10. A 24-year-old male patient has questions regarding variant B.1.1.7. Which of the following educational points should be expressed to the patient?

- A. Variant B.1.1.7 does not affect individuals under the age of 25.
 - B. Variant B.1.1.7 emerged independently in South Africa.
 - C. Variant B.1.1.7 spreads easily and quickly when compared to other COVID-19 variants.
 - D. Variant B.1.1.7 has not been detected in the U.S.
-

11. A fellow health care professional has questions regarding hand hygiene. Which of the following informational points of interest should be communicated to the health care professional regarding hand hygiene?

- A. A health care professional should engage in hand hygiene after touching a patient.
 - B. The duration of the entire handwashing procedure with soap and water should last between 10 - 20 seconds.
 - C. The duration of the entire handwashing procedure with soap and water should last between 5 - 10 seconds.
 - D. At the end of a hand hygiene procedure with an alcohol-based formulation, health care professionals should dry their hands with a towel.
-

12. A health care professional has questions regarding hand hygiene with an alcohol-based formulation. Which of the following informational points of interest should be communicated to the health care professional?

- A. The duration of the entire procedure should last between 5 - 10 seconds.
 - B. The duration of the entire procedure should last between 10 - 15 seconds.
 - C. The duration of the entire procedure should last between 15 - 18 seconds.
 - D. The duration of the entire procedure should last between 20 - 30 seconds.
-

13. Which of the following statements is most accurate?

- A. All individuals with signs and symptoms of a respiratory infection (excluding health care professionals) should be instructed to cover the mouth and nose with their hands when coughing or sneezing.
 - B. All individuals with signs and symptoms of a respiratory infection (including health care professionals) should be instructed to cover the mouth and nose with their hands when coughing or sneezing.
 - C. All individuals over the age of 65 with signs and symptoms of a respiratory infection should be instructed to cover the mouth and nose with their hands when coughing or sneezing.
 - D. All individuals with signs and symptoms of a respiratory infection (including health care professionals) should be instructed to cover the mouth and nose with a tissue when coughing or sneezing.
-

14. A health care professional has questions regarding safe injection practices. Which of the following informational points of interest should be communicated to the health care professional?

- A. Health care professionals should always unwrap syringes prior to the time of use for all patients.
- B. Health care professionals should always unwrap syringes prior to the time of use for patients over the age of 65.
- C. Health care professionals may administer medications from the same syringe to multiple patients.

D. Health care professionals should never administer medications from the same syringe to multiple patients.

15. A fellow health care professional has questions regarding contact precautions. Which of the following informational points of interest should be communicated to the health care professional regarding contact precautions?

- A. If contact precautions may be required, patients with symptoms of an active infection should be placed in a private room until it is determined if contact precaution are necessary.
 - B. Signs indicating patients are on contact precautions should not be posted outside patient rooms.
 - C. PPE is not necessary if contact precautions are required for a patient.
 - D. Patient transport and movement should not be limited.
-

16. Which of the following statements regarding airborne precautions is most accurate?

- A. Health care professionals should ensure appropriate patient placement in an airborne infection isolation room (AIIR); if a health care facility does not have an AIIR, health care professionals should place patients in a private room with the door open.
 - B. Health care professionals should ensure appropriate patient placement in an airborne infection isolation room (AIIR); if a health care facility does not have an AIIR, health care professionals should only place patients over the age of 65 in a private room with the door open.
 - C. Health care professionals should ensure appropriate patient placement in an airborne infection isolation room (AIIR); if a health care facility does not have an AIIR, health care professionals should place patients in a private room with the door closed.
 - D. Health care professionals should ensure appropriate patient placement in an airborne infection isolation room (AIIR); if a health care facility does not have an AIIR, health care professionals should only place patients over the age of 65 in a private room with the door closed.
-

17. A 52-year-old female patient requires droplet precautions. How can health care professionals ensure droplet precautions are applied to the patient?

- A. Ensure the patient wears a face shield and goggles at all times.
 - B. Ensure the patient wears a facemask, face shield, and goggles at all times.
 - C. Allow the patient to exit the room for periods of 20 - 90 minutes.
 - D. Limit the transport and movement of patients outside of their rooms to medically-necessary purposes.
-

18. A health care professional has questions regarding the Interim Infection Prevention and Control Recommendations. Which of the following informational points of interest should be communicated to the health care professional?

- A. Facemasks are preferred over cloth face masks for health care professionals.

- B. Cloth masks are preferred over facemasks for health care professionals.
 - C. Facemasks are not considered to be PPE.
 - D. Cloth masks are considered to be PPE.
-

19. An adult individual enters a health care facility. The adult individual is carrying a child under the age of two. A health care professional observes that the individual and the child are not wearing masks. According to the Interim Infection Prevention and Control Recommendations, how should the health care professional proceed?

- A. Place a facemask on the adult individual and the child.
 - B. Place a respirator on the adult individual and a facemask on the child.
 - C. Offer the adult individual and the child a facemask or mask.
 - D. Offer the adult individual a facemask or mask, and note that facemasks and cloth masks should not be placed on young children under age two.
-

20. A health care professional has questions regarding COVID-19 screening. Which of the following informational points of interest should be communicated to the health care professional?

- A. Screening for fever and symptoms of COVID-19 should be incorporated into the daily assessments of all admitted patients.
 - B. Screening for fever and symptoms of COVID-19 should only be incorporated into the daily assessments of patients over the age of 65.
 - C. Screening for fever and symptoms of COVID-19 should only be incorporated into the daily assessments of patients under the age of 18.
 - D. Screening for fever and symptoms of COVID-19 should not be incorporated into the daily assessments of patients.
-